

学校编码: 10384

分类号_____密级_____

学号: 24520111153309

UDC_____

廈門大學

碩 士 學 位 論 文

酞菁红区荧光探针在生物大分子检测中的应用及其用于生物成像的可行性初探

Red Emitting Phthalocyanines Used as Fluorescent Probes
for the Analysis of Biomacromolecules and Preliminary
Investigation of Their Applications in Bioimaging

陈 林

指导教师姓名 : 李东辉 教授

专 业 名 称 : 药 理 学

论文提交日期 : 2014 年 4 月

论文答辩时间 : 2014 年 5 月

学位授予日期 : 2014 年 月

答辩委员会主席:

评 阅 人:

2014 年 5 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

（ ） 1.经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。

（ ☒ ） 2.不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月 日

目录

摘要	IX
Abstract	XI
第一章 前言	1
1.1 红区及近红外荧光探针的应用研究进展	1
1.1.1 红区及近红外荧光探针的简介	1
1.1.2 红区及近红外荧光探针的分类	1
1.1.3 红区及近红外荧光探针的应用	2
1.2 酞菁化合物的简介	6
1.2.1 酞菁化合物的分子结构和光谱特性	6
1.2.2 酞菁化合物的应用研究	8
1.3 课题设想	13
参考文献	15
第二章 四磺基铝酞菁-阳离子表面活性剂缔合体系用于硫酸软骨素的检测	22
第一节 四磺基铝酞菁-十六烷基溴化吡啶缔合体系用于硫酸软骨素的含量测定	22
2.1.1 引言	22
2.1.2 实验部分	22
2.1.2.1 仪器	22
2.1.2.2 试剂	23
2.1.2.3 实验方法	23
2.1.2.4 批量样品的微孔板测定方法	23
2.1.3 结果与讨论	23
2.1.3.1 AlS_4Pc 的分子结构	23

2.1.3.2 阳离子表面活性剂的选择	25
2.1.3.3 酸度的影响	27
2.1.3.4 反应时间和温度的影响	27
2.1.3.5 标准曲线	27
2.1.3.6 共存物质的干扰	28
2.1.3.7 复杂样品（滴眼液）的测定	28
2.1.3.8 批量样品的测定	30
2.1.4 结论	31
参考文献	32
第二节 四磺基铝酞菁-十四烷基二甲基苄基氯化铵缔合体系用于硫酸软骨素的含量测定	34
2.2.1 引言	34
2.2.2 实验部分	34
2.2.2.1 仪器	34
2.2.2.2 试剂	34
2.2.2.3 实验方法	35
2.2.3 结果与讨论	35
2.2.3.1 AlS_4Pc - TDBAC 体系的激发和发射光谱	35
2.2.3.2 阳离子表面活性剂的选择	36
2.2.3.3 酸度的影响	36
2.2.3.4 反应时间和温度的影响	36
2.2.3.5 标准曲线	37
2.2.3.6 共存物质的干扰	37
2.2.3.7 实际样品（CS 滴眼液）的含量测定	38
2.2.4 结论	40
参考文献	41
第三章 阳离子铝酞菁-黏多糖缔合物检测溶菌酶	43
3.1 引言	43

3.2 实验部分	44
3.2.1 仪器	45
3.2.2 试剂	45
3.2.3 实验方法	45
3.3 结果与讨论	45
3.3.1 TTMAAIPc 的分子结构	45
3.3.2 反应机理探讨	46
3.3.3 酸度的影响	48
3.3.4 缓冲体系的选择	49
3.3.5 反应时间的影响	50
3.3.6 温度的影响	50
3.3.7 标准曲线	51
3.3.8 共存物质的干扰	51
3.3.9 实际样品的测定	52
3.4 结论	54
参考文献	55
第四章 酞菁红区荧光探针用于指纹成像初探	58
4.1 引言	58
4.2 实验部分	58
4.2.1 仪器	58
4.2.2 试剂	58
4.2.3 实验方法	59
4.3 结果和讨论	60
4.3.1 指印的固定	60
4.3.2 染料浓度的考察	61
4.3.3 染料的选择	61
4.3.4 溶剂的选择	61
4.3.5 染色时间的考察	62
4.3.6 讨论	62

参考文献	64
第五章 酞菁红区荧光染料结合显微注射技术用于细胞成像初探 ..	65
5.1 引言	65
5.2 实验部分	65
5.2.1 仪器	65
5.2.2 试剂	65
5.2.3 实验材料	66
5.3 显微注射的操作步骤	66
5.3.1 操作用针的制备	66
5.3.2 注射样品的灌装	67
5.3.3 琼脂糖固定垫的制备	67
5.3.4 细胞的准备	67
5.3.5 注射	68
5.3.6 观察结果	69
5.4 讨论	71
参考文献	72
附录：硕士期间发表论文	73
致谢	75

Contents

Abstract in Chinese	IX
Abstract in English	XI
Chapter 1 Introduction	1
1.1 Research progress of red and near red-region fluorescent probes	1
1.1.1 Introduction of red and near-red region fluorescent probe	1
1.1.2 Classification of near-red region fluorescent probes	1
1.1.3 The application of near-red region fluorescent probes	2
1.2 Introduction of phthalocyanines compounds	6
1.2.1 Molecular structures and spectral characteristics of phthalocyanine compounds	6
1.2.2 The application of phthalocyanines compounds	8
1.3 Thesis Design	13
References	15
Chapter 2 Determination of chondroitin sulfate based on shifting the association equilibrium between tetrasulphonated aluminium phthalocyanine and cationic surfactants	22
2.1 AlS₄Pc-CPB association system for the determination of chondroitin sulfate	22
2.1.1 Introduction	22
2.1.2 Experimental	22
2.1.2.1 Apparatus	22
2.1.2.2 Reagents	23
2.1.2.3 Experimental	23
2.1.2.4 Determination of batch testing	23

2.1.3 Results and discussion	23
2.1.3.1 Molecular structure and spectral characteristic of AlS ₄ Pc	23
2.1.3.2 The selection of cationic surfactants	25
2.1.3.3 Effect of pH	27
2.1.3.4 Effect of reaction time and temperature	27
2.1.3.5 Calibration curve	27
2.1.3.6 Interference	28
2.1.3.7 Analysis of complex samples (eye drops)	28
2.1.3.8 Analysis of batch testing	30
2.1.4 Conclusions	31
References	32
2.2 AlS₄Pc-TDBAC association system for the determination of chondroitin sulfate	34
2.2.1 Introduction	34
2.2.2 Experimental	34
2.2.2.1 Apparatus	34
2.2.2.2 Reagents	34
2.2.2.3 Experimental	35
2.2.3 Results and discussion	35
2.2.3.1 Molecular structure and spectral characteristic of AlS ₄ Pc	35
2.2.3.2 The selection of cationic surfactants	36
2.2.3.3 Effect of pH	36
2.2.3.4 Effect of reaction time and temperature	36
2.2.3.5 Calibration curve	37
2.2.3.6 Interference	37
2.2.3.7 Analysis of practical samples	38
2.2.4 Conclusions	40
References	41

Chapter 3 A novel method for determination of lysozyme based on fluorescence recovery of cationic aluminiumphthalocyanine-mucopoly saccharides association complex used as fluorogenic substrate emitting at red-region	43
3.1 Introduction	43
3.2 Experimental	44
3.2.1 Apparatus	45
3.2.2 Reagents	45
3.2.3 Experimental	45
3.3 Results and discussion	45
3.3.1 Molecular structure and spectral characteristic of TTMAAlPc	45
3.3.2 Discussion of reaction mechanism	46
3.3.3 Effect of pH	48
3.3.4 Effect of buffer system	49
3.3.5 Effect of reaction time	50
3.3.6 Effect of temperature	50
3.3.7 Calibration curve	51
3.3.8 Interference	51
3.3.9 Analysis of practical samples	52
3.4 Conclusions	54
References	55
Chapter 4 Preliminary investigation on the imaging of fingerprints using fluorescent phthalocyanines as stains	58
4.1 Introduction	58
4.2 Experimental	58
4.2.1 Apparatus	58
4.2.2 Reagents	58

4.2.3 Experimental	59
4.3 Results and discussion	60
4.3.1 Fixation of fingerprints	60
4.3.2 Effect of the concentration of phthalocyanine dyes	61
4.3.3 Select of dyes	61
4.3.4 Selection of solvents	61
4.3.5 Effect of staining time	62
4.3.6 Discussion	62
References	64
Chapter 5 Preliminary investigation on cell imaging using fluorescent phthalocyanine as stain	65
5.1 Introduction	65
5.2 Experimental	65
5.2.1 Apparatus	65
5.2.2 Reagents	65
5.2.3 Materials	66
5.3 The procedure of microinjection	66
5.3.1 Preparation of needles for microinjection	66
5.3.2 Filling-in the entry needles with dyes	67
5.3.3 Preparation of agarose fixed pad	67
5.3.4 Preparation of the cells	67
5.3.5 Microinject	68
5.3.6 Results and discussion	69
5.4 Discussion	71
References	72
Appendix	73
Acknowledgements	75

摘要

由于在长波区域具有吸收或发射特性的天然和人工合成的物质极少,因而在进行生物、生化和临床等复杂试样的测定时,使用长波发射荧光试剂可有效避开背景荧光和散射光的干扰,且光漂白作用小,与传统的荧光试剂相比具有很大优越性。能发荧光的酞菁化合物即是一类红区发射的荧光试剂。本论文的工作围绕酞菁红区荧光探针在生物大分子分析及成像检测中的应用而展开,共分五章。

第一章:首先对红区及近红外荧光探针的应用进展进行了简要介绍。由于酞菁化合物作为红区荧光试剂的研究构成了本文的主要部分,本章还对酞菁化合物的分子结构和光谱特性以及它在生物模拟酶、光动力治疗、生化分析领域的应用做了介绍。

第二章:本章建立了简便、快速测定硫酸软骨素的荧光增强分析法。通过荧光光谱筛选实验发现,具有共轭结构的阳离子头部和长碳链尾部的阳离子表面活性剂可几乎完全猝灭四磺基铝酞菁(Tetrasulfonated Aluminum Phthalocyanine, AlS_4Pc)的荧光。而在带有磺基阴离子的硫酸软骨素(Chondroitin Sulfate, CS)存在下,体系荧光显著恢复。以 AlS_4Pc -阳离子表面活性剂离子缔合物为红区荧光探针,考察了其对 CS 的荧光恢复响应行为,发现离子缔合物的荧光恢复程度与 CS 浓度存在良好的相关性,据此实现了复杂样品中 CS 简便、准确的测定。本章工作分为两部分:由于前期工作筛选出的猝灭效果较好的阳离子表面活性剂有两种,由这两种猝灭剂参与的体系的灵敏度和线性范围不同,各具特色,因此对两个体系分别进行了考察,并建立了两种 CS 的荧光增强测定法。

第三章:本章建立了简便、快速测定溶菌酶的荧光增强分析法。带有强阴离子的黏多糖肝素(Heparin, HP)可诱导阳离子铝酞菁[Tetra(trimethyammionio) Aluminum Phthalocynine, $TTMAAlPc$]聚集而导致荧光显著猝灭。由于溶菌酶对黏多糖具有催化降解作用,因而可水解 HP 为小分子片段,从而破坏 $TTMAAlPc$ -HP 的聚集缔合平衡,使 $TTMAAlPc$ 被释放,体系荧光将因之恢复。结合荧光光谱与荧光各向异性技术对反应机理进行了探讨。据此建立了溶菌酶测定新方法,并实现了复杂样品中的溶菌酶含量的准确测定。

第四章：探讨了红区荧光染料酞菁在指纹成像观测中的应用。考察了几种酞菁染料对潜在指印的成像效果，其中 $\text{Al}(\text{SO}_2\text{Cl})_4\text{TSP}$ 对于玻片上油指印的染色效果最为理想，进一步的对染料溶剂，浓度，染色方法和时间进行了考察和优化，成像结果令人满意。

第五章：为研究红区荧光染料酞菁在细胞中是否有特异性定位，为此首先通过摸索建立了显微注射技术，直接将荧光酞菁染料注射进细胞中，体外培养后观察染料与细胞是否有结合。初步研究的结果表明荧光酞菁化合物具有成为活细胞成像新型荧光探针的潜力。

关键词：酞菁；硫酸软骨素；溶菌酶；含量测定；生物成像

Abstract

Long wavelength emitting fluorescent probes can effectively avoid the interference of background fluorescence and scattered light, as compared with the conventional fluorescent reagents, because the natural and synthetic materials have little absorption or emission in long-wavelength region. Phthalocyanine compounds include one class of fluorescent reagents emitted in red region. This work focus on the applications of red emitting phthalocyanine fluorescent probes on the analysis of biomacromolecules and imaging detection, it is divided into five chapters.

In chapter 1, the developments of fluorescent probes in red and near-infrared region were reviewed, followed by the introduction of phthalocynine compounds about their molecular structure, characteristic of molecular spectra, methods for preparation and the main application areas of these compounds.

In chapter 2, this work aims at developing a novel method for rapid determination of chondroitin sulfate. The fluorescence of tetrasulphonated aluminum phthalocyanine (AlS_4Pc), a anionic metal phthalocyanine, was quenched dramatically by cationic surfactant in which contains a positively-charged head with a conjugated structure and a long carbon chain as tail through the formation of a almost non-fluorescent association complex. It was found that the ion-association complex ($\text{AlS}_4\text{Pc-CPB}$) emitted strong fluorescence in the presence of chondroitin sulfate, due to the ability of chondroitin sulfate to shift the association equilibrium of the ion-association complex that led to the release of AlS_4Pc , thus resulting in an increase in the fluorescence of AlS_4Pc . Based on the above-mentioned phenomenon, a novel method for quantitative determination of chondroitin sulfate in practical samples was developed using the ion-association phthalocyanine complex as a fluorescent probe emitting at red-region. Two anionic surfactants quenching the fluorescence of AlS_4Pc with high efficiency were screened. It was found that different sensitivity and linear range for the determination of chondroitin sulfate were obtained using the two surfactants as quenchers. Therefore, two determination

systems were investigated separately.

In chapter 3, the main idea of this part of work is to develop a novel method for rapid determination of lysozyme. It was found that the fluorescence of the cationic aluminum phthalocyanine, a red-region fluorescence probe, was extremely quenched in acidic media in the presence of low concentrations of anionic mucopolysaccharide heparin (HP) bearing anionic sulfonic acid groups, due to induced aggregation. The almost non-fluorescent substrate was degraded into small molecule fragments by the hydrolysis of lysozyme, hence, the phthalocyanine molecules aggregated in HP were released, resulted in significant fluorescence recovery of the reaction system. This phenomenon bases the principle of the proposed method. The reaction mechanism was discussed employing fluorescence spectroscopy and steady-state fluorescence anisotropy techniques. Factors which affected the determination were investigated. The established method has been used to the analysis of real samples of lysozyme, the results are in well agreement with those determined by a conventional turbidimetric method.

In chapter 4, preliminary investigation on the the application of fluorescent phthalocyanines for the imaging of potent fingerprints are carried out. Seven fluorescent phthalocyanines were screened, and factors which affected the detection were investigated. Among phthalocyanine compounds having been investigated, $\text{Al}(\text{SO}_2\text{Cl})_4\text{TSP}$ showed the most effective results for the imaging of potent fingerprints.

In the final chapter, the possibility of fluorescent phthalocyanines for in vivo imaging of cells was preliminarily investigated. The purpose of this work is to study the location behavior of fluorescent phthalocyanines inside cells. Microinjection technique was established for injecting fluorescent phthalocyanines into cells directly. Preliminary results of the study showed that fluorescent phthalocyanine compounds have the great potential to be new fluorescent probes for cell imaging in vivo.

Keywords: phthalocyanine; chondroitin sulfate; lysozyme; determination; bio-imaging

Degree papers are in the “[Xiamen University Electronic Theses and Dissertations Database](#)”. Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库